



TITLE:

Search for Eulerian Recurrent Lengths by Using Constraint Solvers (Developments of Language, Logic, Algebraic system and Computer Science)

AUTHOR(S):

Jimbo, Shuji

CITATION:

Jimbo, Shuji. Search for Eulerian Recurrent Lengths by Using Constraint Solvers (Developments of Language, Logic, Algebraic system and Computer Science). 数理解析研究所講究録 2017, 2051: 52-56

ISSUE DATE:

2017-10

URL:

<http://hdl.handle.net/2433/237089>

RIGHT:

Search for Eulerian Recurrent Lengths by Using Constraint Solvers

Shuji JIMBO

Graduate School of Natural Science and Technology, Okayama University

jimbo-s@okayama-u.ac.jp

Abstract

The Eulerian recurrent length of a graph G , $e(G)$, is the maximum of the shortest subcycle length of Eulerian circuits of G . Upper and lower bounds on the Eulerian recurrent length of complete graphs was provided by the author as $n - 4 \leq e(K_n) \leq n - 3$ for odd integers $n \geq 15$. In this article, the method of proving the inequality is improved.

KEYWORDS. graph theory, Eulerian circuits, computer experiments, integer programming, Eulerian recurrent length.

1 Introduction

Let $C = v_0 \rightarrow v_1 \rightarrow v_2 \rightarrow \cdots \rightarrow v_{m-1} \rightarrow v_0$ be a circuit of a graph, where m is the length of C . For any integer k , we regard v_k as the vertex $v_{k \bmod m}$ in C for convenience of discussion, where $k \bmod m$ is the minimum nonnegative integer of the form $k - qm$ with integer q . We call a subwalk $W = v_i \rightarrow v_{i+1} \rightarrow \cdots \rightarrow v_j$ in C a subcycle if $v_i = v_j$ and W is a cycle of K_n . The shortest subcycle length of C is denoted by $s(C)$. Let G be a Eulerian graph. We call $e(G) = \max\{s(C) \mid C \text{ is an Eulerian circuit of } G\}$ the Eulerian recurrent length (ERL) of G . Let K_n denote the complete graph with n vertices, where n is an odd positive integer with $n \geq 3$. Then, K_n is a Eulerian graph. We abbreviate $e(K_n)$ as $e(n)$ in this paper.

We have proved the following fact. Let k be an arbitrary integer greater than 330. Let G denote a given four-regular Eulerian graph. Then, the problem to determine whether $e(G) \geq k$ or not is NP-complete[1]. We also determined the ERL of complete bipartite graphs as follows. Let $K_{m,n}$ denote the complete bipartite graph of vertex classes with m and n vertices. Let m and n be even integers with $0 < n < m$. Then, $e(K_{n,n}) = 2n - 4$ and $e(K_{m,n}) = 2n$ hold[4][2]. We also determine the ERL of complete graphs K_n , namely $e(n)$, for particular small integers n by computer verification. Equation $e(3) = e(5) = 3$ holds, and, for any $n \in \{7, 9, 11, 13\}$, equation $e(n) = n - 3$ holds. For complete graphs K_n with odd integer $n \geq 15$, we have had the following upper and lower bounds on $e(n)$:

$$n - 4 \leq e(K_n) \leq n - 3.$$

The left inequality above can be obtained by showing a construction method of a Eulerian circuit C of K_n with a shortest subcycle of length $n - 4$. The construction method is based

on a decomposition of $E(K_n)$, the edge set of K_n , into Hamiltonian cycles. The Eulerian circuit C can be obtained by aligning the Hamiltonian cycles in appropriate order, and by joining them[2]. We will describe the method of proving the right inequality above by solving integer programming problems in the previous article[2]. Then, we will give an improvement of the proof by a modification to the integer programming problem.

2 Main arguments

For a circuit C of length m and an integer i , $C(i)$ denotes the vertex on position $i \bmod m$ in C , and is called the i -th vertex on C . Hence, $C = C(0) \rightarrow C(1) \rightarrow \cdots \rightarrow C(m-1) \rightarrow C(0)$ holds, and any two edges $C(i)C(i+1)$ and $C(j)C(j+1)$ are different if $i < j < i+m$. For any integer i , $N_C(i)$ denotes the unique integer k such that

$$i < k \leq i+m, \quad C(k) = C(i), \quad \text{and} \quad C(j) \neq C(i) \quad \text{for any } j \text{ with } i < j < k.$$

Hence, for any integer i , $N_C^{-1}(i)$ denotes the unique integer k such that $N_C(k) = i$.

Suppose that C is an Eulerian circuit of a complete graph K_n . An edge $e = C(i)C(i+1)$ is negative if either

$$N_C^{-1}(i+1) < N_C^{-1}(i) \quad \text{and} \quad N_C(i) < N_C(i+1)$$

or

$$N_C^{-1}(i) < N_C^{-1}(i+1) \quad \text{and} \quad N_C(i+1) < N_C(i)$$

holds, and positive otherwise. A quadruple (i, j, k, l) of integers is a (non-contact) position reversal, or PR, on C , if $i < j < k = N_C(j) < l = N_C(i)$ holds. The following proposition readily follows from the definitions above.

Proposition 1 *For any Eulerian circuit C of K_n with $n \geq 7$, the number of PR's on C is not less than that of negative edges on C , where two PR's (i, j, k, l) and (i', j', k', l') are regarded as identical if $i \equiv i' \pmod{m}$, $j \equiv j' \pmod{m}$, $k \equiv k' \pmod{m}$, and $l \equiv l' \pmod{m}$ hold.*

For a position reversal $r = (i, j, k, l)$ on C , i and l are called the position reversal head and the position reversal tail of r , respectively.

Let $n \geq 7$ be an odd integer. Let m denote $n(n-1)/2$, the number of edges of K_n . We assume that there exists an Eulerian circuit C with $s(C) = n-2$ of K_n to prove that $e(n) \leq n-3$ by contradiction. We have the following two lemmas by tedious arguments[3].

Lemma 1 *For any integer i ,*

$$n-2 \leq N_C(i) - i \leq n+3$$

holds.

Lemma 2 *For any integer i , if i is a position reversal head on C , then the following equations hold:*

$$\begin{aligned} N_C(i) - i &= n+3, \\ N_C(i+1) - (i+1) &= n-2, \quad \text{and} \\ N_C(i+2) - (i+2) &= n-1. \end{aligned}$$

Furthermore, for any integer i , if i is a position reversal tail on C , then the following equations hold:

$$\begin{aligned} i - N_C^{-1}(i) &= n + 3, \\ (i - 1) - N_C^{-1}(i - 1) &= n - 2, \text{ and} \\ (i - 2) - N_C^{-1}(i - 2) &= n - 1. \end{aligned}$$

For an integer p and a positive integer μ , $S_\mu(p)$ denotes the set $\{p, p+1, \dots, p+\mu-1\}$. Then, $A_\mu^C(i)$, $B_\mu^C(i)$, and $C_\mu^C(i)$ denote the number of left positions of negative edges on C , position reversal heads on C , and position reversal tails on C in $S_\mu(i)$, respectively. The following lemma follows from Proposition 1 immediately.

Lemma 3 *Let n be an integer greater than or equal to 7. Let C be an Eulerian circuit of K_n . Let m denote $n(n-1)/2$, the length of C . Let μ be a positive integer less than m . Then, the following two inequalities hold:*

$$\sum_{i=0}^{m-1} (B_\mu^C(i) - A_\mu^C(i)) \geq 0, \text{ and} \quad (1)$$

$$\sum_{i=0}^{m-1} (B_\mu^C(i) + C_\mu^C(i) - 2A_\mu^C(i)) \geq 0. \quad (2)$$

Corollary 1 *Let n be an integer greater than or equal to 7. Let C be an Eulerian circuit of K_n . Let m denote $n(n-1)/2$, the length of C . Let μ be a positive integer less than m . Then, there is an integer $p \in \{0, 1, 2, \dots, m-1\}$ such that*

$$B_\mu^C(p) - A_\mu^C(p) \geq 0. \quad (3)$$

There also is $q \in \{0, 1, 2, \dots, m-1\}$ such that

$$B_\mu^C(q) + C_\mu^C(q) - 2A_\mu^C(q) \geq 0. \quad (4)$$

Suppose that an integer p and a positive integer μ are given. Let $x(i) = (p + i - 2) - N_C^{-1}(p + i - 2)$ and $y(i) = N_C(p + i - 2) - (p + i - 2)$ for each $i \in \{0, 1, 2, \dots, \mu + 3\}$. Then, the following conditions must hold by definition:

- (a) $\forall i \in \{0, 1, \dots, \mu + 3\}, \forall j \in \{0, 1, \dots, \mu + 3\}, i \neq j \Rightarrow (i - x(i) \neq j - x(j) \wedge i + y(i) \neq j + y(j))$,
- (b) $\forall i \in \{0, 1, \dots, \mu + 2\}, |x(i + 1) - x(i) - 1| \neq 1 \wedge |y(i) - y(i + 1) - 1| \neq 1$,
- (c) $\forall i \in \{0, 1, \dots, \mu + 3\}, \forall j \in \{0, 1, \dots, \mu + 3\}, i \neq j \Rightarrow (|(i - x(i)) - (j - x(j))| \neq 1 \vee |(i + y(i)) - (j + y(j))| \neq 1)$,
- (d1) $\forall i \in \{0, 1, \dots, \mu - 2\}, \exists j \in \{0, 1, 2, 3, 4, 5\}, x(i + j) = j$, and
- (d2) $\forall i \in \{0, 1, \dots, \mu - 2\}, \exists j \in \{0, 1, 2, 3, 4, 5\}, y(i + j) = 5 - j$.

Let X and Y denote $(x(0), x(1), \dots, x(\mu+3))$ and $(y(0), y(1), \dots, y(\mu+3))$, respectively. Let $M_\mu(X, Y)$ denote the number of left positions of negative edges in $\{p, p+1, \dots, p+\mu-1\}$. Notice that, for each $i \in \{0, 1, 2, \dots, \mu+3\}$, $p+i-2$ is a left position of a negative edge on C , if and only if the following condition holds:

$$(x(i) < x(i+1) \wedge y(i) < y(i+1)) \vee (x(i) > x(i+1) \wedge y(i) > y(i+1)).$$

Let $R_\mu(X, Y)$ denote the number of position reversal heads in $\{p, p+1, \dots, p+\mu-1\}$. Notice that, for each $i \in \{0, 1, 2, \dots, \mu+3\}$, $p+i-2$ is a position reversal head on C , if and only if the following condition holds:

$$y(i) = 5 \wedge y(i+1) = 0 \wedge y(i+2) = 1.$$

Let $R'_\mu(X, Y)$ denote the number of position reversal tails in $\{p, p+1, \dots, p+\mu-1\}$. Notice that, for each $i \in \{0, 1, 2, \dots, \mu+3\}$, $p+i-2$ is a position reversal tail on C , if and only if the following condition holds:

$$x(i) = 5 \wedge x(i-1) = 0 \wedge x(i-2) = 1.$$

The problem finding the maximum value of $R_\mu(X, Y) - M_\mu(X, Y)$ under the conditions (a), (b), (c), (d1) and (d2) can be formulated as an integer programming problem. By solving the problem, we have:

- There is no position i on C such that $R_7(X, Y) > M_7(X, Y)$.
- For any position i on C , $M_7(X, Y) \geq 2$ holds.

Those conditions are too tight for C . Additional arguments therefore lead us to contradiction. Thus, we have proved that there is no Eulerian circuit C of K_n with $s(C) = n-2$ for any odd integer $n \geq 15$ [3]. However, a modification to the integer programming problem saves the additional arguments above as described in the next paragraph.

The problem finding the maximum value of $R_\mu(X, Y) + R'_\mu(X, Y) - 2M_\mu(X, Y)$ under the conditions (a), (b), (c), (d1), and (d2) can be formulated as an integer programming problem. By solving the problem, we have:

- There is no position i on C s.t. $R_7(X, Y) + R'_7(X, Y) \geq M_7(X, Y)$.

It directly follows from the statement above that there is no Eulerian circuit C of K_n with $s(C) = n-2$. Thus, the proof have been simplified. We consider that complexity of the integer programming problem is substituted for a part of arguments in the previous proof.

3 Concluding Remarks

We has described an improvement of the previous proof of the upper bound on the Eulerian recurrent length of complete graphs K_n for odd integers $n \geq 7$. The improvement has been brought by a modification to the integer programming problem used for deriving a contradiction.

The gap between the upper and lower bound on the Eulerian recurrent length of K_n , $e(n)$, for odd $k \geq 15$ is still remains as $n-4 \leq e(n) \leq n-3$. We conjecture that

$e(n) = n - 4$ holds for any odd integer $n \geq 15$. Currently, we try to verify that $e(15) = 11$ by computer. However, we estimate the number of primitive checks on graphs required by computer experiments for the verification according to naive methods at about 3^{45} or more. The computer experiments, therefore, should result in failure in the current computation environment.

Acknowledgment

We appreciate kindly discussion by the members in the RIMS workshop. This work was supported by JSPS KAKENHI Grant Number 15K00018.

References

- [1] Shuji Jimbo. The np-completeness of eulerian recurrent length for 4-regular eulerian graphs. In *Artificial Intelligence with Applications in Engineering and Technology (ICAJET), 2014 4th International Conference on*, pages 155–159. IEEE, 2014.
- [2] Shuji Jimbo. On the eulerian recurrent lengths of complete bipartite graphs and complete graphs. In *IOP Conference Series: Materials Science and Engineering*, volume 58(1), page 012019. IOP Publishing, 2014.
- [3] Shuji Jimbo. Improvement of the upper bound on the eulerian recurrent lengths of complete graphs. In *The Proceedings of the 9th Hungarian-Japanese Symposium on Discrete Mathematics and Its Applications*. IOP Publishing, 6 2015.
- [4] Shuji Jimbo, Yasuaki Oshie, and Kosaburo Hashiguchi. The NP-completeness of EULERIAN RECURRENT LENGTH. *RIMS Kôkyûroku*, 1437:107–115, 2005.